



TricaJus

User Guide  
Penerangan

说明手册

The continuous page are shown the possible reactions after taking TricaJus

Mukasurat berikut menunjukkan reaksi yang mungkin dialami selepas minum TricaJus

内页内容提及可能性的调理反应

Tricajus Tricajus Tricajus Tricajus Tricajus Tricajus Tricajus Tricajus

[www.macamix.myhealth2u.com](http://www.macamix.myhealth2u.com)

Body Reactions / Reaksi Badan / 反应	Possible Reasons / Sebab - sebab / 病因	Solution / Cara makan / 处理方法
<ul style="list-style-type: none"> <li><b>Slight fever or body feels heaty.</b> Demam ringan atau badan berasa panas. 发烧或发热。</li> </ul>	<ul style="list-style-type: none"> <li><b>Improved circulation with promotes body metabolism to release body heat and hence increase body temperature.</b> Peningkatan peredaran darah menyebabkan metabolisme badan meningkat. Haba dibebaskan untuk mengimbangkan suhu badan. 促进血液循环提高新陈代谢, 产生热量, 增加体温。</li> </ul>	<ul style="list-style-type: none"> <li><b>Take in the morning. Avoid taking at night. Drink more water.</b> Minum pada waktu pagi, elakkan dari meminumnya pada waktu malam. minum banyak air. 早上服用, 避免晚上服用。和大量水份。</li> </ul>
<ul style="list-style-type: none"> <li><b>Fever, feeling alert, excited &amp; can't sleep.</b> Demam, terangsang, bertenaga atau terjaga pada waktu malam. 发烧, 微亢奋, 精神, 难眠。</li> </ul>	<ul style="list-style-type: none"> <li><b>Burning of excessive fats that converts heat in energy.</b> Penukaran lemak yang berlebihan ke dalam bentuk tenaga. 身体把多余的脂肪转化为热能。</li> </ul>	<ul style="list-style-type: none"> <li><b>Take in the morning. Avoid taking at night. Drink more water.</b> Minum pada waktu pagi, elakkan dari meminumnya pada waktu malam. minum banyak air. 早上服用, 避免晚上服用。喝大量水。</li> </ul>
<ul style="list-style-type: none"> <li><b>Feces colour turns dark.</b> Kumuhan hijau kehitam-hitaman. 排泄物呈黑色。</li> </ul>	<ul style="list-style-type: none"> <li><b>Detoxification resulting from action by fibre.</b> Fiber bertindak membuang sisa toksin yang terkumpul di dalam kolon. 纤维帮助清除积在肠内已久的残余物。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake. Avoid heavy foods. Drink more water.</b> Minum setiap hari, kurangkan makanan yang terlalu berat, minum banyak air. 继续每天服用, 饮食清淡。喝大量水。</li> </ul>
<ul style="list-style-type: none"> <li><b>Flatulence, uneasy chest experience, nausea.</b> Berkentut, sakit dada, sedu dan loya. 频密放屁, 胸口发闷, 打呃, 作呕。</li> </ul>	<ul style="list-style-type: none"> <li><b>Malfunction of digestive system.</b> Sistem penghadaman lemah atau mengalami kekembungan. 肠胃不好。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Sleepy &amp; tiredness.</b> Mengantuk, tangan atau kaki berasa letih. 爱睡, 手脚疲倦。</li> </ul>	<ul style="list-style-type: none"> <li><b>Acidic blood malfunction of liver, over exhausted, too much drugs &amp; chemicals.</b> Badan berasid dan fungsi hati lemah, tertekan, terlalu letih atau pengambilan ubat berlebihan. 酸性体质, 肝脏机能衰退。压力, 疲劳或服药太多所致。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake. Drink more water. Less meat &amp; more rest.</b> Minum setiap hari, minum banyak air, kurangkan makan daging dan banyak berehat. 每天服用, 喝大量水。少吃肉类, 多休息。</li> </ul>
<ul style="list-style-type: none"> <li><b>Dizziness, weak &amp; hungry.</b> Pening, lemah, selalu berasa lapar. 晕眩, 虚弱, 饥饿。</li> </ul>	<ul style="list-style-type: none"> <li><b>Lower blood sugar level or anemia. Fat will be burn off &amp; blood sugar will be lowered.</b> Menghadap hipoglisemia atau anemia. Paras gula darah turun untuk membakar lemak yang berlebihan. 低血糖或贫血。脂肪燃烧时, 血糖会降低。</li> </ul>	<ul style="list-style-type: none"> <li><b>Take after meal. Avoid consume on an empty stomach.</b> Minum selepas makan. Jangan minum dengan perut kosong. 餐后服用, 避免空腹服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Diarrhoea.</b> Cirit - bitit. 下痢, 腹泻。</li> </ul>	<ul style="list-style-type: none"> <li><b>Malfunction of digestive system, fibre help for detoxification.</b> Fiber sedang membersihkan toksin badan. 纤维帮助排毒。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Constipation, bleeding in passing motion.</b> Sembelit, Kumuhan berdarah. 便秘, 大便出血或血丝现像。</li> </ul>	<ul style="list-style-type: none"> <li><b>Piles, insufficient fibre, water or mineral.</b> Penyakit buasir, kekurangan fiber, air atau mineral di dalam badan. 痔疮, 缺乏纤维素, 水份或矿物质。</li> </ul>	<ul style="list-style-type: none"> <li><b>Take in the morning. Avoid taking at night. Drink more water.</b> Minum pada waktu pagi, elakkan dari meminumnya pada waktu malam. minum banyak air. 早上服用, 避免晚上服用。喝大量水。</li> </ul>
<ul style="list-style-type: none"> <li><b>Pimples, rash, sweating.</b> Jerawat, bintik - bintik merah, berpeluh. 青春痘, 红斑, 出汗。</li> </ul>	<ul style="list-style-type: none"> <li><b>Poor function of liver, excessive toxin in our body.</b> Jantung lemah, terdapat toksin berlebihan di dalam badan. 肝脏机能衰退, 毒素太多。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Menstruation disorder.</b> Haid yang tidak tepat. 月经不规则。</li> </ul>	<ul style="list-style-type: none"> <li><b>Imbalance of hormone in body.</b> Masalah ketidakseimbangan hormon badan. 荷尔蒙失调。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Runny nose, asthma, coughs expectoration.</b> Hidung berair, asma, berkahak. 流鼻水, 气喘, 多咳, 多痰。</li> </ul>	<ul style="list-style-type: none"> <li><b>Smoker, weak respiratory function.</b> Penghisap rokok, fungsi respirasi lemah. 吸烟者, 呼吸系统衰退。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Slight edema on face &amp; foot, weight gain.</b> Muka dan kaki bengkak, berat badan naik. 脸肿, 脚肿, 体重上升。</li> </ul>	<ul style="list-style-type: none"> <li><b>Poor function of kidney &amp; heart.</b> Fungsi buah pinggang, jantung dan hati yang lemah. 肾及心脏机能衰退。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Paralysis of hand &amp; leg.</b> Kaki tangan rasa kebas, kejang. 手脚麻痹, 抽筋。</li> </ul>	<ul style="list-style-type: none"> <li><b>Bad circulation, insufficient calcium.</b> Peredaran darah lemah, kekurangan kalsium di dalam badan. 血液循环不良, 缺钙。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>
<ul style="list-style-type: none"> <li><b>Inflammation of old injuries.</b> Kawasan yang dijangkit / luka lebih sakit. 患处加剧疼痛。</li> </ul>	<ul style="list-style-type: none"> <li><b>Improved circulation, old injury is being cured.</b> Peredaran darah sedang ditingkatkan untuk memperbaiki kawasan yang jangkit / luka. 促进血液调节, 修补病患处。</li> </ul>	<ul style="list-style-type: none"> <li><b>Continue intake.</b> Minum berterusan. 继续每天服用。</li> </ul>